



IVTScrip™ mRNA-Human ADAMTS4, (Cap 0, 5-Methyl-CTP & Pseudo-UTP, 30 nt-poly(A))

Cat. No.: GTTS-WK10703MR

This product is for research use only and is not intended for diagnostic use.

PRODUCT INFORMATION

Product overview

This product GTTS-WK10703MR is a type of mRNA having 120 nt poly(A) tail and modified with Cap 0 & 5-Methyl-CTP & Pseudo-UTP. It encodes the ADAMTS4 protein. This product can be used in Oligodendrocyte-related researches.

Specifications

Modified bases	5-Methyl-CTP & Pseudo-UTP
5' Cap	Cap 0
Species	Human
RefSeq	NM_001320336.3
Applications	Gene therapy research
Format	Powder
Quantity	100 µg
Purification	Chromatography

SPECIFICATIONS

Modified bases	5-Methyl-CTP & Pseudo-UTP
5' Cap	Cap 0
Species	Human
RefSeq	NM_001320336.3
Applications	Gene therapy research
Format	Powder
Quantity	100 µg
Purification	Chromatography

GENE INFORMATION

Alternative Names	ADMP-1; ADAMTS-2; ADAMTS-4
Description	This gene encodes a member of the ADAMTS (a disintegrin and metalloproteinase with thrombospondin motifs) protein family. Members of this family share several distinct protein modules, including a propeptide region, a metalloproteinase domain, a disintegrin-like domain, and a thrombospondin type 1 (TS) motif. Individual members of this family differ in the number of C-terminal TS motifs, and some have unique C-terminal domains. The enzyme encoded by this gene lacks a C-terminal TS motif. The encoded preproprotein is proteolytically processed to generate the mature protease. This protease is responsible for the degradation of aggrecan, a major proteoglycan of cartilage, and brevican, a brain-specific extracellular matrix protein. The expression of this gene is upregulated in arthritic disease and this may contribute to disease progression through the degradation of aggrecan. Alternative splicing results in multiple transcript variants, at least one of which encodes an isoform that is proteolytically processed. [provided by RefSeq, Feb 2016]