



IVTScrip™ mRNA-Human ADH5, (Cap 0, 5-Methyl-CTP & Pseudo-UTP, 30 nt-poly(A))

Cat. No.: GTTS-WK10749MR

This product is for research use only and is not intended for diagnostic use.

PRODUCT INFORMATION

Product overview

This product GTTS-WK10749MR is a type of mRNA having 120 nt poly(A) tail and modified with Cap 0 & 5-Methyl-CTP & Pseudo-UTP. It encodes the ADH5 protein. This product can be used in Natural killer T (NKT) cell-related researches.

Specifications

Modified bases	5-Methyl-CTP & Pseudo-UTP
5' Cap	Cap 0
Species	Human
RefSeq	NM_000671.4
Applications	Gene therapy research
Format	Powder
Quantity	100 µg
Purification	Chromatography

SPECIFICATIONS

Modified bases	5-Methyl-CTP & Pseudo-UTP
5' Cap	Cap 0
Species	Human
RefSeq	NM_000671.4
Applications	Gene therapy research
Format	Powder
Quantity	100 µg
Purification	Chromatography

GENE INFORMATION

Alternative Names	FDH; ADHX; ADH-3; AMEDS; BMFS7; FALDH; GSNOR; GSH-FDH; HEL-S-60p
Description	This gene encodes a member of the alcohol dehydrogenase family. Members of this family metabolize a wide variety of substrates, including ethanol, retinol, other aliphatic alcohols, hydroxysteroids, and lipid peroxidation products. The encoded protein forms a homodimer. It has virtually no activity for ethanol oxidation, but exhibits high activity for oxidation of long-chain primary alcohols and for oxidation of S-hydroxymethyl-glutathione, a spontaneous adduct between formaldehyde and glutathione. This enzyme is an important component of cellular metabolism for the elimination of formaldehyde, a potent irritant and sensitizing agent that causes lacrymation, rhinitis, pharyngitis, and contact dermatitis. The human genome contains several non-transcribed pseudogenes related to this gene. [provided by RefSeq, Oct 2008]