



## IVTScrip™ mRNA-Human ATP2A1, (Cap 1, 5-Methyl-CTP & Pseudo-UTP, 120 nt-poly(A))

Cat. No.: GTTS-WK26989MR

This product is for research use only and is not intended for diagnostic use.

### PRODUCT INFORMATION

#### Product overview

This product GTTS-WK26989MR is a type of mRNA having 120 nt poly(A) tail and modified with Cap 1 & 5-Methyl-CTP & Pseudo-UTP. It encodes the ATP2A1 protein. This product can be used in Enterocyte progenitor cell-related researches.

#### Specifications

<b>Modified bases</b>	5-Methyl-CTP & Pseudo-UTP
<b>5' Cap</b>	Cap 1
<b>Species</b>	Human
<b>RefSeq</b>	NM_001286075.2
<b>Applications</b>	Gene therapy research
<b>Format</b>	Powder
<b>Quantity</b>	100 µg
<b>Purification</b>	Chromatography

### SPECIFICATIONS

<b>Modified bases</b>	5-Methyl-CTP & Pseudo-UTP
<b>5' Cap</b>	Cap 1
<b>Species</b>	Human
<b>RefSeq</b>	NM_001286075.2
<b>Applications</b>	Gene therapy research
<b>Format</b>	Powder
<b>Quantity</b>	100 µg
<b>Purification</b>	Chromatography

## GENE INFORMATION

<b>Alternative Names</b>	ATP2A; SERCA1
<b>Description</b>	This gene encodes one of the SERCA Ca(2+)-ATPases, which are intracellular pumps located in the sarcoplasmic or endoplasmic reticula of muscle cells. This enzyme catalyzes the hydrolysis of ATP coupled with the translocation of calcium from the cytosol to the sarcoplasmic reticulum lumen, and is involved in muscular excitation and contraction. Mutations in this gene cause some autosomal recessive forms of Brody disease, characterized by increasing impairment of muscular relaxation during exercise. Alternative splicing results in three transcript variants encoding different isoforms. [provided by RefSeq, Oct 2013]