



IVTScrip™ mRNA-Human ADAMTS2, (Cap 0, 5-Methoxy-UTP, 120 nt-poly(A))

Cat. No.: GTTS-WK8002MR

This product is for research use only and is not intended for diagnostic use.

PRODUCT INFORMATION

Product overview

This product GTTS-WK8002MR is a type of mRNA having 120 nt poly(A) tail and modified with Cap 0 & 5-Methoxy-UTP. It encodes the ADAMTS2 protein. This product can be used in SLC16A7+ cell-related researches.

Specifications

Modified bases	5-Methoxy-UTP
5' Cap	Cap 0
Species	Human
RefSeq	NM_014244.5
Applications	Gene therapy research
Format	Powder
Quantity	100 µg
Purification	Chromatography

SPECIFICATIONS

Modified bases	5-Methoxy-UTP
5' Cap	Cap 0
Species	Human
RefSeq	NM_014244.5
Applications	Gene therapy research
Format	Powder
Quantity	100 µg
Purification	Chromatography

GENE INFORMATION

Alternative Names	NPI; PNPI; PCINP; PCPNI; PCI-NP; PC I-NP; ADAM-TS2; ADAMTS-2; ADAMTS-3; EDSDERMS
Description	This gene encodes a member of the ADAMTS (a disintegrin and metalloproteinase with thrombospondin motifs) protein family. Members of the family share several distinct protein modules, including a propeptide region, a metalloproteinase domain, a disintegrin-like domain, and a thrombospondin type 1 (TS) motif. Individual members of this family differ in the number of C-terminal TS motifs, and some have unique C-terminal domains. The encoded preproprotein is proteolytically processed to generate the mature procollagen N-proteinase. This proteinase excises the N-propeptide of the fibrillar procollagens types I-III and type V. Mutations in this gene cause Ehlers-Danlos syndrome type VIIC, a recessively inherited connective-tissue disorder. Alternative splicing results in multiple transcript variants, at least one of which encodes an isoform that is proteolytically processed. [provided by RefSeq, Feb 2016]