



## IVTScrip™ mRNA-Human ACE2, (Cap 0, 5-Methoxy-UTP, 30 nt-poly(A))

Cat. No.: GTTS-WK9722MR

This product is for research use only and is not intended for diagnostic use.

### PRODUCT INFORMATION

#### Product overview

This product GTTS-WK9722MR is a type of mRNA having 120 nt poly(A) tail and modified with Cap 0 & 5-Methoxy-UTP. It encodes the ACE2 protein. This product can be used in Oogenesis phase fetal germ cell-related researches.

#### Specifications

<b>Modified bases</b>	5-Methoxy-UTP
<b>5' Cap</b>	Cap 0
<b>Species</b>	Human
<b>RefSeq</b>	NM_001371415.1
<b>Applications</b>	Gene therapy research
<b>Format</b>	Powder
<b>Quantity</b>	100 µg
<b>Purification</b>	Chromatography

### SPECIFICATIONS

<b>Modified bases</b>	5-Methoxy-UTP
<b>5' Cap</b>	Cap 0
<b>Species</b>	Human
<b>RefSeq</b>	NM_001371415.1
<b>Applications</b>	Gene therapy research
<b>Format</b>	Powder
<b>Quantity</b>	100 µg
<b>Purification</b>	Chromatography

## GENE INFORMATION

<b>Alternative Names</b>	ACEH
<b>Description</b>	<p>The protein encoded by this gene belongs to the angiotensin-converting enzyme family of dipeptidyl carboxydipeptidases and has considerable homology to human angiotensin 1 converting enzyme. This secreted protein catalyzes the cleavage of angiotensin I into angiotensin 1-9, and angiotensin II into the vasodilator angiotensin 1-7. ACE2 is known to be expressed in various human organs, and its organ- and cell-specific expression suggests that it may play a role in the regulation of cardiovascular and renal function, as well as fertility. In addition, the encoded protein is a functional receptor for the spike glycoprotein of the human coronavirus HCoV-NL63 and the human severe acute respiratory syndrome coronaviruses, SARS-CoV and SARS-CoV-2, the latter is the causative agent of coronavirus disease-2019 (COVID-19). Multiple splice variants have been found for this gene and the dACE2 (or MIRb-ACE2) splice variant has been found to be interferon inducible. [provided by RefSeq, Nov 2020]</p>