



## IVTScrip™ mRNA-Human ACP2, (Cap 0, 5-Methoxy-UTP, 30 nt-poly(A))

Cat. No.: GTTS-WK9739MR

This product is for research use only and is not intended for diagnostic use.

### PRODUCT INFORMATION

#### Product overview

This product GTTS-WK9739MR is a type of mRNA having 120 nt poly(A) tail and modified with Cap 0 & 5-Methoxy-UTP. It encodes the ACP2 protein. This product can be used in Gonadal endothelial cell-related researches.

#### Specifications

<b>Modified bases</b>	5-Methoxy-UTP
<b>5' Cap</b>	Cap 0
<b>Species</b>	Human
<b>RefSeq</b>	NM_001302489.2
<b>Applications</b>	Gene therapy research
<b>Format</b>	Powder
<b>Quantity</b>	100 µg
<b>Purification</b>	Chromatography

### SPECIFICATIONS

<b>Modified bases</b>	5-Methoxy-UTP
<b>5' Cap</b>	Cap 0
<b>Species</b>	Human
<b>RefSeq</b>	NM_001302489.2
<b>Applications</b>	Gene therapy research
<b>Format</b>	Powder
<b>Quantity</b>	100 µg
<b>Purification</b>	Chromatography

## GENE INFORMATION

<b>Alternative Names</b>	LAP
<b>Description</b>	<p>The protein encoded by this gene belongs to the histidine acid phosphatase family, which hydrolyze orthophosphoric monoesters to alcohol and phosphate. This protein is localized to the lysosomal membrane, and is chemically and genetically distinct from the red cell acid phosphatase. Mice lacking this gene showed multiple defects, including bone structure alterations, lysosomal storage defects, and an increased tendency towards seizures. An enzymatically-inactive allele of this gene in mice showed severe growth retardation, hair-follicle abnormalities, and an ataxia-like phenotype. Alternatively spliced transcript variants have been found for this gene. A C-terminally extended isoform is also predicted to be produced by the use of an alternative in-frame translation termination codon via a stop codon readthrough mechanism. [provided by RefSeq, Oct 2017]</p>